PATIENT HEALTH QUESTIONNAIRE (PHQ-9)

ID #:	DATE:			
Over the last 2 weeks, how often have you been				
bothered by any of the following problems? (use "✓" to indicate your answer)	Not at all	Several days	More than half the days	Nearly every day
Little interest or pleasure in doing things	0 -	1	2	3
2. Feeling down, depressed, or hopeless	0	1	2	3
3. Trouble falling or staying asleep, or sleeping too much	0	1	2	3
4. Feeling tired or having little energy	0	1	2	3
5. Poor appetite or overeating	0	1	1	3
6. Feeling bad about yourself—or that you are a failure or have let yourself or your family down	0	1	2	3
7. Trouble concentrating on things, such as reading the newspaper or watching television	0	1	2 · 2 · 2	3
8. Moving or speaking so slowly that other people could have noticed. Or the opposite — being so figety or restless that you have been moving around a lot more than usual	0	1	2	3
9. Thoughts that you would be better off dead, or of hurting yourself	0	1 - 1 - 1 - 1 - 1 - 1 - 1 - 1	2	3
	add columns	5	+	+
(Healthcare professional: For interpretation of TOT please refer to accompanying scoring card).	<i>TAL</i> , TOTAL			
10. If you checked off any problems, how difficult		Not d	fficult at all	(Section 1)
have these problems made it for you to do		Some	what difficult	Weinsk
your work, take care of things at home, or get	Very difficult			
along with other people?	Extremely difficult			

Copyright $\mathbb C$ 1999 Pfizer Inc. All rights reserved. Reproduced with permission. PRIME-MD $\mathbb C$ is a trademark of Pfizer Inc. A2663B 10-04-2005

Generalized Anxiety Disorder 7-item (GAD-7) scale

Over the last 2 weeks, how often have you been bothered by the following problems?	Not at all sure	Several days	Over half the days	Nearly every day
1. Feeling nervous, anxious, or on edge	0	1	2	3
2. Not being able to stop or control worrying	0	1	2	3
3. Worrying too much about different things	0	1	2	3
4. Trouble relaxing	0	1	2	3
5. Being so restless that it's hard to sit still	0	1	2	3
6. Becoming easily annoyed or irritable	0	1	2	3
7. Feeling afraid as if something awful might happen	0	1	2	3
Add the score for each column	+	+	+	
Total Score (add your column scores) =				

If you checked off any problems, how difficult have these made it for you to do your work, take care of things at home, or get along with other people?

Not difficult at all	
Somewhat difficult	
Very difficult	
Extremely difficult	

Scoring

Scores of 5, 10, and 15 are taken as the cut-off points for mild, moderate and severe anxiety, respectively. When used as a screening tool, further evaluation is recommended when the score is 10 or greater.

Using the threshold score of 10, the GAD-7 has a sensitivity of 89% and a specificity of 82% for GAD. It is moderately good at screening three other common anxiety disorders - panic disorder (sensitivity 74%, specificity 81%), social anxiety disorder (sensitivity 72%, specificity 80%) and post-traumatic stress disorder (sensitivity 66%, specificity 81%).

Source: Spitzer RL, Kroenke K, Williams JBW, Lowe B. A brief measure for assessing generalized anxiety disorder. *Arch Inern Med.* 2006;166:1092-1097.